


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C07D 207/16, A61K 31/401, A61P 3/10</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/34241 (43) International Publication Date: 15 June 2000 (15.06.00)</p>
<p>(21) International Application Number: PCT/EP99/09708 (22) International Filing Date: 9 December 1999 (09.12.99) (30) Priority Data: 09/209,068 10 December 1998 (10.12.98) US (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventor; and (75) Inventor/Applicant (for US only): VILLHAUER, Edwin, Bernard [US/US]; 20 Dorothy Drive, Morristown, NJ 07960 (US). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent &amp; Trademark Dept., CH-4002 Basel (CH).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: N-SUBSTITUTED 2-CYANOPYRROLIDINES</p> <div style="text-align: center;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>The present invention relates to a compound of formula (I), wherein R is substituted adamantyl; and n is 0 to 3; in free form or in acid addition salt form. Compounds of formula (I) inhibit DPP-IV (dipeptidyl-peptidase-IV) activity. They are therefore indicated for use as pharmaceuticals in inhibiting DPP-IV and in the treatment of conditions mediated by DPP-IV, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, osteoporosis and further conditions of impaired glucose tolerance.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## N-SUBSTITUTED 2-CYANOPYRROLIDINES

The present invention provides new dipeptidyl peptidase-IV (DPP-IV) inhibitors which are effective in treating conditions mediated by DPP-IV. More recently, it was discovered that DPP-IV is responsible for inactivating glucagon-like peptide-1 (GLP-1). Since GLP-1 is a major stimulator of pancreatic insulin secretion and has direct beneficial effects on glucose disposal, DPP-IV inhibition appears to represent an attractive approach for treating conditions such as non-insulin-dependent diabetes mellitus (NIDDM).

The instant invention relates to novel N-(substituted glycy)-2-cyanopyrrolidines of formula I:



wherein

R is substituted adamantyl; and

n is 0 to 3; in free form or in acid addition salt form.

The compounds of formula I can exist in free form or in acid addition salt form. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolating or purifying the compounds of this invention. Although the preferred acid addition salts are the hydrochlorides, salts of methanesulfonic, sulfuric, phosphoric, citric, lactic and acetic acid may also be utilized.

The compounds of the invention may exist in the form of optically active isomers or diastereoisomers and can be separated and recovered by conventional techniques, such as chromatography.

Listed below are definitions of various terms used to describe this invention. These definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either individually or as part of a larger group.

The term "alkyl" refers to straight or branched chain hydrocarbon groups having 1 to 10 carbon atoms, preferably 1 to 7 carbon atoms, most preferably 1 to 5 carbon atoms. Exemplary alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl and the like.

The term "alkanoyl" refers to alkyl-C(O)-.

The term "substituted adamantyl" refers to adamantyl, i.e. 1- or 2-adamantyl, substituted by one or more, for example two, substituents selected from alkyl, -OR<sub>1</sub> or -NR<sub>2</sub>R<sub>3</sub>; where R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are independently hydrogen, alkyl, (C<sub>1</sub>-C<sub>8</sub>-alkanoyl), carbamyl, or -CO-NR<sub>4</sub>R<sub>5</sub>; where R<sub>4</sub> and R<sub>5</sub> are independently alkyl, unsubstituted or substituted aryl and where one of R<sub>4</sub> and R<sub>5</sub> additionally is hydrogen or R<sub>4</sub> and R<sub>5</sub> together represent C<sub>2</sub>-C<sub>7</sub>alkylene;.

The term "aryl" preferably represents phenyl. Substituted phenyl preferably is phenyl substituted by one or more, e.g. two, substituents selected from e.g. alkyl, alkoxy, halogen and trifluoromethyl.


The term "alkoxy" refers to alkyl-O-.

The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.

The term "alkylene" refers to a straight chain bridge of 2 to 7 carbon atoms, preferably of 3 to 6 carbon atoms, most preferably 5 carbon atoms.


A preferred group of compounds of the invention is the compounds of formula I wherein the substituent on the adamantyl is bonded on a bridgehead or a methylene adjacent to a bridgehead. Compounds of formula I wherein the glycyl-2-cyanopyrrolidine moiety is bonded to a bridgehead, the R' substituent on the adamantyl is preferably 3-hydroxy. Compounds of formula I wherein the glycyl-2-cyanopyrrolidine moiety is bonded at a methylene adjacent to a bridgehead, the R' substituent on the adamantyl is preferably 5-hydroxy.

The present invention especially relates to a compound of formulae (I A) or (I B)



wherein R' represents hydroxy, C<sub>1</sub>-C<sub>7</sub>alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkanoyloxy, or R<sub>5</sub>R<sub>4</sub>N-CO-O-, where R<sub>4</sub> and R<sub>5</sub> independently are C<sub>1</sub>-C<sub>7</sub>alkyl or phenyl which is unsubstituted or substituted by a substituent selected from C<sub>1</sub>-C<sub>7</sub>alkyl, C<sub>1</sub>-C<sub>7</sub>alkoxy, halogen and trifluoromethyl and where R<sub>4</sub> additionally is hydrogen; or R<sub>4</sub> and R<sub>5</sub> together represent C<sub>3</sub>-C<sub>6</sub>alkylene; and R'' represents hydrogen; or R' and R'' independently represent C<sub>1</sub>-C<sub>7</sub>alkyl; in free form or in form of a pharmaceutically acceptable acid addition salt.

The compounds of the invention may be prepared e.g. by a process which comprises coupling a reactive (2-cyanopyrrolidino)carbonylmethylene compound with an appropriate substituted amine; more particularly, for the preparation of the compounds of formula I, it comprises reacting a compound of formula II



wherein Y is a reactive group (preferably a halogen such as bromine, chlorine or iodine) with a compound of formula III




wherein R is as defined above, and recovering the resultant compound of formula I in free form or in acid addition salt form.

The process of the invention may be effected in conventional manner. For example, the compound of formula II is reacted with 1 to 3 equivalents, preferably 3 equivalents of a primary amine of formula III. The reaction is conveniently conducted in the presence of an inert, organic solvent, such as methylene chloride or a cyclic ether such as tetrahydrofuran. The temperature preferably is of from about 0° to about 35°C, preferably between about 0° and about 25°C.

The compounds of the invention may be isolated from the reaction mixture and purified in conventional manner, e.g. by chromatography.

The starting materials may also be prepared in conventional manner. The compounds of formula II may be prepared by the following two-step reaction scheme:



Step 1 involves the reaction of the pyrrolidine of formula IV with a slight molar excess of a haloacetylhalide such as bromoacetyl bromide or chloroacetyl chloride and a base such as potassium carbonate or triethylamine. The reaction conveniently is conducted in the presence of an inert, organic solvent, such as tetrahydrofuran or a chlorinated, aliphatic hydrocarbon such as methylene chloride, at a temperature of from about 0° to about 25°C, preferably at a temperature between about 0° and about 15°C.

Step 2 concerns the dehydration of the compound of formula V, prepared in Step 1, with 1 to 2 equivalents of trifluoroacetic anhydride (TFAA). The dehydration preferably is conducted in the presence of an inert, organic solvent such as tetrahydrofuran or a chlorinated, aliphatic hydrocarbon such as methylene chloride, at a temperature of from about 0° to about 25°C, preferably at a temperature between about 0° and about 15°C.

Insofar as its preparation is not particularly described herein, a compound used as starting material is known or may be prepared from known compounds in known manner or analogously to known methods or analogously to methods described in the Example.

For example, the primary amine compounds of formula III are known and may be prepared by procedures documented in the literature, for example, Khim. -Farm. Zh. (1986), 20(7), 810 -15.

Finally, compounds of the invention are either obtained in the free form, or as a salt thereof if salt forming groups are present.

Compounds of the invention having basic groups can be converted into acid addition salts, especially pharmaceutically acceptable acid addition salts. These are formed, for example, with inorganic acids, such as mineral acids, for example sulfuric acid, a phosphoric or hydrohalic acid, or with organic carboxylic acids. Preferred are salts formed with hydrochloric acid.

In view of the close relationship between the free compounds and the compounds in the form of their salts, whenever a compound is referred to in this context, a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization.

The instant invention also includes pharmaceutical compositions, for example, useful in inhibiting DPP-IV, comprising a pharmaceutically acceptable carrier or diluent and a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable acid addition salt thereof.

In still another embodiment, the instant invention provides a method of inhibiting DPP-IV comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable acid addition salt thereof.

In a further embodiment, the instant invention provides a method of treating conditions mediated by DPP-IV inhibition comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula I above, or a pharmaceutically acceptable acid addition salt thereof.

The present invention also relates to the use of a compound according to the instant invention or a pharmaceutically acceptable salt thereof e.g. for the manufacture of a medicament for the prevention or treatment of diseases or conditions associated with elevated levels of DPP-IV.

As indicated above, all of the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, are useful in inhibiting DPP-IV. The ability of the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, to inhibit DPP-IV may be demonstrated employing the Caco-2 DPP-IV Assay which measures the ability of test compounds to inhibit DPP-IV activity from human colonic carcinoma cell extracts. The human colonic carcinoma cell line Caco-2 was obtained from the American Type Culture Collection (ATCC HTB 37). Differentiation of the cells to induce DPP-IV expression was accomplished as described by Reisher, et al. in an article entitled "Increased expression of intestinal cell line Caco-2" in Proc. Natl. Acad. Sci., Vol. 90, pgs. 5757-5761 (1993). Cell extract is prepared from cells solubilized in 10mM Tris HCl, 0.15 M NaCl, 0.04 t.i.u. aprotinin, 0.5% nonidet-P40, pH 8.0, which is centrifuged at 35,000 g for 30 min. at 4°C. to remove cell debris. The assay is conducted by adding 20 µg solubilized

Caco-2 protein, diluted to a final volume of 125  $\mu$ l in assay buffer (25 mM Tris HCl pH 7.4, 140mM NaCl, 10 mM KCl, 1% bovine serum albumin) to microtiter plate wells. After a 60 min. incubation at room temperature, the reaction is initiated by adding 25 $\mu$ l of 1 mM substrate (H-Alanine-Proline-pNA; pNA is *p*-nitroaniline). The reaction is carried out at room temperature for 10 minutes after which time a 19  $\mu$ l volume of 25% glacial acetic acid is added to stop the reaction. Test compounds are typically added as 30  $\mu$ l additions and the assay buffer volume is reduced to 95  $\mu$ l. A standard curve of free *p*-nitroaniline is generated using 0-500  $\mu$ M solutions of free pNA in assay buffer. The curve generated is linear and is used for interpolation of substrate consumption (catalytic activity in nmoles substrate cleaved /min). The endpoint is determined by measuring absorbance at 405 nm in a Molecular Devices UV Max microtiter plate reader.

The potency of the test compounds as DPP-IV inhibitors, expressed as IC<sub>50</sub>, is calculated from 8-point, dose-response curves using a 4-parameter logistic function.

The following IC<sub>50</sub> was obtained:

Compound	Caco-2 DPP-IV (nM)
Ex. 1	3.5 $\pm$ 1.5
Ex. 4	8

The ability of the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, to inhibit DPP-IV may also be demonstrated by measuring the effects of test compounds on DPP-IV activity in human and rat plasma employing a modified version of the assay described by Kubota, et al. in an article entitled "Involvement of dipeptidylpeptidase IV in an *in vivo* immune response" in Clin. Exp. Immunol., Vol. 89, pgs. 192-197 (1992). Briefly, 5  $\mu$ l of plasma are added to 96-well flat-bottom microtiter plates (Falcon), followed by the addition of 5  $\mu$ l of 80 mM MgCl<sub>2</sub> in incubation buffer (25 mMHEPES, 140 mM NaCl, 1% RIA-grade BSA, pH 7.8). After a 60 min. incubation at room temperature, the reaction is initiated by the addition of 10  $\mu$ l of incubation buffer containing 0.1 mM substrate (H-Glycine-Proline-AMC; AMC is 7-amino-4-methylcoumarin). The plates are covered with aluminum foil (or kept in the dark) and incubated at room temperature for 20 min. After the 20 min. reaction, fluorescence is measured using a CytoFluor 2350 fluorimeter (Excitation 380 nm Emission 460nm; sensitivity setting 4). Test compounds are



typically added as 2  $\mu$ l additions and the assay buffer volume is reduced to 13  $\mu$ l. A fluorescence-concentration curve of free AMC is generated using 0-50  $\mu$ M solutions of AMC in assay buffer. The curve generated is linear and is used for interpolation of substrate consumption (catalytic activity in nmoles substrate cleaved/min). As with the previous assay, the potency of the test compounds as DPP-IV inhibitors, expressed as  $IC_{50}$ , is calculated from 8-point, dose-response curves using a 4 parameter logistic function.

The following  $IC_{50}$  was obtained:

Compound	human plasma DPP-IV (nM)	rat plasma DPP-IV (nM)
Ex. 1	$2.7 \pm 0.1$	$2.3 \pm 0.1$
Ex. 8	6	12

In view of their ability to inhibit DPP-IV, the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, are useful in treating conditions mediated by DPP-IV inhibition. Based on the above and findings in the literature, it is expected that the compounds disclosed herein are useful in the treatment of conditions such as non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation and calcitonin-osteoporosis. In addition, based on the roles of glucagon-like peptides (such as GLP-1 and GLP-2) and their association with DPP-IV inhibition, it is expected that the compounds disclosed herein are useful for example, to produce a sedative or anxiolytic effect, or to attenuate post-surgical catabolic changes and hormonal responses to stress, or to reduce mortality and morbidity after myocardial infarction, or in the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels.

More specifically, for example, the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, improve early insulin response to an oral glucose challenge and, therefore, are useful in treating non-insulin-dependent diabetes mellitus. The ability of the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, to improve early insulin response to an oral glucose challenge may be measured in insulin resistant rats according to the following method:

Male Sprague-Dawley rats that had been fed a high fat diet (saturated fat = 57% calories) for 2-3 weeks were fasted for approximately 2 hours on the day of testing, divided

into groups of 8-10, and dosed orally with 10  $\mu\text{mol/kg}$  of the test compounds in CMC. An oral glucose bolus of 1 g/kg was administered 30 minutes after the test compound directly into the stomach of the test animals. Blood samples, obtained at various timepoints from chronic jugular vein catheters, were analyzed for plasma glucose and immunoreactive insulin (IRI) concentrations, and plasma DPP-IV activity. Plasma insulin levels were assayed by a double antibody radioimmunoassay (RIA) method using a specific anti-rat insulin antibody from Linco Research (St. Louis, MO). The RIA has a lower limit of detection of 0.5  $\mu\text{U/mL}$  with intra- and inter-assay variations of less than 5%. Data are expressed as % increase of the mean of the control animals. Upon oral administration, each of the compounds tested amplified the early insulin response which led to an improvement in glucose tolerance in the insulin resistant test animals. The following results were obtained:

Compound	Increase of Insulin Response at 10 $\mu\text{mol/kg}$
Ex. 1	64%

The precise dosage of the compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, to be employed for treating conditions mediated by DPP-IV inhibition depends upon several factors, including the host, the nature and the severity of the condition being treated, the mode of administration and the particular compound employed. However, in general, conditions mediated by DPP-IV inhibition are effectively treated when a compound of formula I, or a corresponding pharmaceutically acceptable acid addition salt, is administered enterally, e.g., orally, or parenterally, e.g., intravenously, preferably orally, at a daily dosage of 0.002-5, preferably 0.02-2.5 mg/kg body weight or, for most larger primates, a daily dosage of 0.1-250, preferably 1-100 mg. A typical oral dosage unit is 0.01-0.75 mg/kg, one to three times a day. Usually, a small dose is administered initially and the dosage is gradually increased until the optimal dosage for the host under treatment is determined. The upper limit of dosage is that imposed by side effects and can be determined by trial for the host being treated.

The compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, may be combined with one or more pharmaceutically acceptable carriers

and, optionally, one or more other conventional pharmaceutical adjuvants and administered enterally, e.g., orally, in the form of tablets, capsules, caplets, etc. or parenterally, e.g., intravenously, in the form of sterile injectable solutions or suspensions. The enteral and parenteral compositions may be prepared by conventional means.


The compounds of formula I, and their corresponding pharmaceutically acceptable acid addition salts, may be formulated into enteral and parenteral pharmaceutical compositions containing an amount of the active substance that is effective for treating conditions mediated by DPP-IV inhibition, such compositions in unit dosage form and such compositions comprising a pharmaceutically acceptable carrier.

The compounds of formula I (including those of each of the subscopes thereof and each of the examples) may be administered in enantiomerically pure form (e.g., ee >98%, preferably >99%) or together with the R enantiomer, e.g., in racemic form. The above dosage ranges are based on the compounds of formula I (excluding the amount of the R enantiomer).

The following examples show representative compounds encompassed by this invention and their synthesis. However, it should be clearly understood that they are for purposes of illustration only.

#### Example 1

Pyrrolidine, 1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-, (S)



#### A. 1-Aminoadamantane-3-ol:

Slight modifications to the synthesis found in Khim. -Farm. Zh. (1986), 20(7), 810 - 15, may be used.

To a rapidly stirred, clear and colorless, ice-water chilled mixture of concentrated sulfuric acid 96% (210 mL; 3,943 mmol) and 65% nitric acid (21.0 mL; 217.0 mmol) is added 21.0g (112.0 mmol) of 1-adamantylamine HCl (99%), in small portions over 30 minutes.

Upon adamantylamine hydrochloride addition, slight bubbling occurs and the reaction is slightly exothermic. This bubbling, yellow solution is stirred at ice-water temperature for about 2 hours and then at room temperature for 30 hours. This clear, light yellow reaction is then poured into about 100g of ice and the resulting solution is clear green-blue.

The solution is placed in an ice-water bath and allowed to stir for 30 minutes. Approximately 550g of 89% pure KOH (8.74 mol) is then added in small portions over 45 minutes. During this addition, the reaction is exothermic; reaching 80°C and producing copious amounts of brown NO<sub>2</sub> gas. By the end of the addition, the reaction is thick with white solids (both product and salts). The resulting white paste is then poured onto a buchner funnel/celite pad and washed with 1.2 L of CH<sub>2</sub>Cl<sub>2</sub>. The CH<sub>2</sub>Cl<sub>2</sub> layer is then extracted from the water layer and dried over Na<sub>2</sub>SO<sub>4</sub>. The solution is then filtered and concentrated (rotovap/pump) to provide 1-aminoadamantane-3-ol as a white solid.

#### B. 1-Chloroacetyl-2-cyanopyrrolidine

To a mechanically stirred solution of 20.0g (180.0mmol) of chloroacetylchloride and 97g (0.70mmol) of potassium carbonate in 150mL of tetrahydrofuran is added a solution of L-prolinamide 20.0g (180.0mmol) in 500 mL of tetrahydrofuran in a dropwise fashion over 45 minutes. This reaction is then mechanically stirred for an additional two hours at room temperature. The reaction is then filtered to remove potassium salts and the filtrate is dried over Na<sub>2</sub>SO<sub>4</sub>. The Na<sub>2</sub>SO<sub>4</sub> is then removed via filtration and to this colorless filtrate is added trifluoroacetic anhydride (25.0mL, 0.180mmol) in one portion. The reaction is then magnetically stirred for 1 hour at room temperature and the resulting clear yellow/orange solution is concentrated via rotovap. The excess trifluoroacetic anhydride is removed by adding ethyl acetate to the concentrated oil and reconcentrating via rotovap. This removing operation is performed three times.



The resulting oil is partitioned between ethyl acetate and water. The product is then extracted into the ethyl acetate and the aqueous layer is then washed twice with ethyl acetate. The combined organic layers are then washed successively with water and brine dried over magnesium sulfate, filtered and concentrated to obtain 1-chloroacetyl-2-cyanopyrrolidine as a yellow solid.




#### C. Pyrrolidine, 1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-, (S)




To a heterogeneous solution of the title A compound (1-aminoadamantane-3-ol (5.80g, 34.7mmol) in  $\text{CH}_2\text{Cl}_2$  (68.0mL) is added 9.6g (69mmol) of  $\text{K}_2\text{CO}_3$ . This heterogeneous mixture is then cooled in an ice-water bath and a solution of 3.0g (17mmol) of the title B compound (1-chloroacetyl-2-cyanopyrrolidine) dissolved in 25.0mL of  $\text{CH}_2\text{Cl}_2$  is added dropwise over a period of 30 minutes. The resulting mixture is stirred for 2 hours at  $0^\circ\text{C}$  and at room temperature for 6 days. The reaction is then concentrated to obtain a yellow pasty material which is purified on silica gel employing a SIMS/Biotage Flash chromatography system and a 7% solution of methanol in methylene chloride as the eluent to yield the title compound in free base form as a white crystalline solid (melting point  $138^\circ\text{C}$ - $140^\circ\text{C}$ ,  $^{13}\text{C}$ NMR (ppm) = 119.59).




### Examples 2 to 12

The following compounds are prepared analogous to the method of Example 1 (especially Step C):

Example	Structure	M.P. [ $^\circ\text{C}$ ]
2	 <p>Pyrrolidine, 1-[(3,5-dimethyl-1-adamantyl)amino]acetyl-2-cyano-, (S)-</p>	103-105 (HCl)
3	 <p>Pyrrolidine, 1-[(3-ethyl-1-adamantyl)amino]acetyl-2-cyano-, (S)-</p>	212-214 (HCl)

4	 <p>Pyrrolidine, 1-[[[(3-methoxy-1-adamantyl)amino]acetyl]-2-cyano-, (S)-</p>	92-94 (HCl)
5	 <p>Chiral</p> <p>Pyrrolidine, 1-[[[3-[[[(t-butylamino)carbonyl]oxy]-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	210-212 (HCl)
6	 <p>Chiral</p> <p>Pyrrolidine, 1-[[[3-[[[(4-methoxyphenyl)amino]carbonyl]oxy]-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	212-214 (HCl)

7	 <p>Pyrrolidine, 1-[[[(3-[[[(phenylamino)carbonyl]oxy]-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	205-207 (HCl)
8	 <p>Pyrrolidine, 1-[[[(5-hydroxy-2-adamantyl)amino]-acetyl]-2-cyano-, (S)-</p>	[ <sup>13</sup> C NMR (CN group): 121.56 (ppm)] (HCl)
9	 <p>Pyrrolidine, 1-[[[(3-acetyloxy-1-adamantyl)amino]-acetyl]-2-cyano-, (S)-</p>	[ <sup>13</sup> C NMR (CN group): 118.54 (ppm)]

10	 <p>Pyrrolidine, 1-[[[3-[[[(diisopropyl)amino]carbonyl]oxy]-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	Chiral 148-150 (HCl)
11	 <p>Pyrrolidine, 1-[[[3-[[[(cyclohexyl)amino]carbonyl]oxy]-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	Chiral 155-157 (HCl)
12	 <p>Pyrrolidine, 1-[[[3-ethoxy-1-adamantyl]amino]acetyl]-2-cyano-, (S)-</p>	Chiral [ <sup>13</sup> C NMR (CN group): 119.31 (ppm)] (HCl)

(HCl) = as hydrochloride

All HCl salts of final products are prepared by passing HCl gas through a 0.1 Molar solution of the free base in tetrahydrofuran until solution is clearly acidic followed by removal of the solvent (rotovap/pump).



The amino-adamantane starting materials are known in the literature or can be prepared as follows:

The manufacture of 3,5-dimethyl-1-adamantylamine is described in J. Med. Chem, 25; 1; 1982; 51-56.

The manufacture of 3-ethyl-1-adamantylamine is described in J. Med. Chem, 25; 1; 1982; 51-56.

3-Methoxy-1-adamantylamine can be prepared as follows:

To a stirred, ice-water chilled suspension of potassium hydride (0.680 gm; 5.95 mmol) in 15.0 ml of tetrahydrofuran is added a mixture of 1-aminoadamantane-3-ol (1.00g; 5.95 mmol) and 15.0 ml of tetrahydrofuran dropwise over 30 minutes. The resulting mixture is then stirred for an additional 30 minutes and iodomethane (0.370 ml; 5.95 mmol) is then added dropwise over one minute. The resulting opaque white reaction is then stirred at room temperature for 18 hours. The mixture is then diluted with 50 ml of methylene chloride and filtered to remove the inorganic impurities. The filtrate is then concentrated and purified on silica gel employing a SIMS/Blotage apparatus and 19% methanol and 1% ammonium hydroxide in methylene chloride as eluent to yield 3-methoxy-1-adamantylamine as an opaque oil.

Synthesis of 3-[(tertbutylamino)carbonyloxy]-1-aminoadamantane:

To a mixture of 1-aminoadamantane-3-ol (5.00 g; 30.0 mmol) and potassium carbonate (6.20 g; 45 mmol) in 150 ml of tetrahydrofuran is added benzylchloroformate (4.70 g, 33.0 mmol) in dropwise fashion over a 10 minute period. The mixture is then stirred at room temperature for 2 h and then partitioned between ethyl acetate and water. The product is then extracted into the ethyl acetate and the aqueous layer is washed twice with ethyl acetate (100 ml). The combined organic layers are then washed successively with 100 ml of aqueous 2 N sodium hydroxide, water and brine, dried over sodium sulfate, filtered and concentrated (rotovap/pump) to provide 1-benzylcarbamoyladamantane-3-ol as a white solid in 85% yield.

To a clear solution of 1-benzylcarbamoyladamantane-3-ol (1.00 g; 3.32 mmol) and tert-butylisocyanate (380 µl, 3.32 mmol) in 30 ml of methylene chloride is syringe-added trimethylsilyl chloride (20.0 µl, 0.17 mmol). This reaction is then stirred at room temperature

for 18 hours, concentrated (rotovap) and purified on silica gel employing a SIMS/Biotage apparatus and 20% ethyl acetate in hexane as eluent to yield 3-[[[(tertbutylamino)carbonyl]oxy]-1-benzylcarbamoyladamantane as a white solid in quantitative yield.

To a mixture of 3-[[[(tertbutylamino)carbonyl]oxy]-1-benzylcarbamoyladamantane (1.50 g, 3.75 mmol) and 10% palladium on carbon (400 mg) in ethanol (150 ml) in a 1-liter parr hydrogenation flask is added hydrogen (50 psi). This opaque black mixture is then shaken for 24 h. The reaction is then filtered through celite to remove the palladium catalyst and concentrated (rotovap/pump) to provide 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane as a clear oil in 99% yield.

The procedure for the synthesis of 4-[[[(methoxyphenyl)amino]carbonyl]oxy]-1-aminoadamantane is essentially the procedure of 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane except in the second step where an equivalent of 4-methoxyphenyl isocyanate replaces tert-butylisocyanate, 1,2-dichloroethane is used as solvent instead of methylene chloride and the reaction is stirred at 50°C for 18 hours. The final amine intermediate is provided as an oil.

The procedure for the synthesis of 3-[[[(phenylamino)carbonyl]oxy]-1-aminoadamantane is essentially the procedure of 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane except in the second step where an equivalent of phenyl isocyanate replaces the tert-butylisocyanate, 1,2-dichloroethane is used as solvent instead of methylene chloride and the reaction is stirred at 50°C for 18 hours. The final amine intermediate is provided as a clear oil.

The procedure to make 2-aminoadamantane-5-ol is the same as in Example 1 except that the starting material is 2-aminoadamantane instead of 1-aminoadamantane.

The procedure for the synthesis of the nucleophile 3-acetoxy-1-aminoadamantane is essentially the procedure of 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane except for a standard acylation of 1-benzylcarbamoyladamantane-3-ol using 1.2 eq of acetyl chloride, 3.0 eq. of pyridine, 0.1 eq of 4-dimethylaminopyridine and 1,2 dichloroethane which are all stirred at room temperature for 24 hours. The final amine is provided as a thick oil.

The procedure for the synthesis of 3-[[[(diisopropyl)amino]carbonyl]oxy]-1-amino-adamantane is essentially the procedure of 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane except in the second step where an equivalent of diisopropylcarbonyl chloride replaces the tert-butyliisocyanate, 1,2-dichloroethane is used as solvent instead of methylene chloride and the reaction is stirred at 85°C for 18 hours. The final amine intermediate is provided as a gray solid.

The procedure for the synthesis of 3-[[[(cyclohexyl)amino]carbonyl]oxy]-1-aminoadamantane is essentially the procedure of 3-[[[(tertbutylamino)carbonyl]oxy]-1-aminoadamantane except in the second step where an equivalent of cyclohexylisocyanate replaces the tert-butyliisocyanate, 1,2-dichloroethane is used as solvent instead of methylene chloride and the reaction is stirred at 50°C for 18 hours. The final amine intermediate is provided as a thick clear oil.

The procedure to make 3-ethoxy-1-adamantylamine (a clear oil) is the same as for 3-methoxy-1-adamantylamine except that iodoethane (1.3 equivalent) is used instead of iodomethane.

#### Formulation Example:

Tablets, each containing 50 mg of active ingredient, for example, (S)1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-pyrrolidine, can be prepared as follows:


#### Composition (for 10,000 tablets)

Active ingredient	500.0 g
Lactose	500.0 g
Potato starch	352.0 g
Gelatin	8.0 g
Talc	60.0 g
Magnesium stearate	10.0 g
Silica (highly disperse)	20.0 g
Ethanol	q.s.

The active ingredient is mixed with the lactose and 292 g of potato starch, and the mixture is moistened using an alcoholic solution of the gelatin and granulated by means of a sieve. After drying, the remainder of the potato starch, the talc, the magnesium stearate and the highly disperse silica are admixed and the mixture is compressed to give tablets of weight 145.0 mg each and active ingredient content 50.0 mg which, if desired, can be provided with breaking notches for finer adjustment of the dose.

## WHAT IS CLAIMED IS:


1. A compound of formula I:



wherein

R is substituted adamantyl; and  
n is 0 to 3; in free form or in acid addition salt form.

2. A compound according to claim 1 of formula e (I A) or (I B)



wherein R' represents hydroxy, C<sub>1</sub>-C<sub>7</sub>alkoxy, C<sub>1</sub>-C<sub>8</sub>-alkanoyloxy, or R<sub>5</sub>R<sub>4</sub>N-CO-O-, where R<sub>4</sub> and R<sub>5</sub> independently are C<sub>1</sub>-C<sub>7</sub>alkyl or phenyl which is unsubstituted or substituted by a substituent selected from C<sub>1</sub>-C<sub>7</sub>alkyl, C<sub>1</sub>-C<sub>7</sub>alkoxy, halogen and trifluoromethyl and where R<sub>4</sub> additionally is hydrogen; or R<sub>4</sub> and R<sub>5</sub> together represent C<sub>3</sub>-C<sub>8</sub>alkylene; and R'' represents hydrogen; or R' and R'' independently represent C<sub>1</sub>-C<sub>7</sub>alkyl; in free form or in form of a pharmaceutically acceptable acid addition salt.

3. A compound according to claim 1 of formula I selected from the group consisting of:

pyrrolidine, 1-[[[(3,5-dimethyl-1-adamantyl)amino]-acetyl]-2-cyano-, (S)-;

pyrrolidine, 1-[[[3-ethyl-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-methoxy-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[1-butylamino)carbonyl]oxy]-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[4-methoxyphenyl)amino]carbonyl]oxy]-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[phenylamino)carbonyl]oxy]-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[5-hydroxy-2-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[3-acetyloxy-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[diisopropyl)amino]carbonyl]oxy]-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
pyrrolidine, 1-[[[3-[[[cyclohexyl)amino]carbonyl]oxy]-1-adamantyl)amino]acetyl]-2-cyano-, (S)-; and  
Pyrrolidine, 1-[[[3-ethoxy-1-adamantyl)amino]acetyl]-2-cyano-, (S)-;  
or, in each case, a pharmaceutically acceptable acid addition salt thereof.

4. A compound according to claim 1 which is: pyrrolidine, 1-[[[3-hydroxy-1-adamantyl)amino]acetyl]-2-cyano-, (S), or a pharmaceutically acceptable salt thereof.

5. A pharmaceutical composition comprising a compound according to claim 1 in free form or in pharmaceutically acceptable acid addition salt form, together with at least one pharmaceutically acceptable carrier or diluent.

6. Use of a compound according to claim 1 or a pharmaceutically acceptable acid addition salt thereof for the manufacture of a medicament for inhibiting dipeptidyl peptidase-IV; or  
a method of inhibiting dipeptidyl peptidase-IV comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound according to claim 1, or a pharmaceutically acceptable acid addition salt thereof.

7. Use according to claim 6 wherein the medicament is used for treating conditions mediated by dipeptidyl peptidase-IV inhibition, or  
a method of treating conditions mediated by dipeptidyl peptidase-IV inhibition comprising administering to a mammal in need of such treatment a therapeutically effective amount of

a compound according to Claim 1, or a pharmaceutically acceptable acid addition salt thereof.

8. Use according to claim 7 wherein the medicament is used for the treatment of non-insulin-dependent diabetes mellitus; or  
the method according to claim 7 wherein the condition treated is non-insulin-dependent diabetes mellitus.

9. Use according to claim 7 wherein the medicament is used for the treatment of obesity; or  
the method according to claim 7 wherein the condition treated is obesity.

# INTERNATIONAL SEARCH REPORT

Inter. Appl. No.  
PCT/EP 99/09708

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D207/16 A61K31/401 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 19998 A (NOVARTIS AG) 14 May 1998 (1998-05-14) the whole document	1,5
A	WO 95 15309 A (FERRING B.V.) 8 June 1995 (1995-06-08) page 11 -page 31	1,5
A	SIMON J. COUTTS: "Structure-Activity Relationships of Boronic acid inhibitors of Dipeptidyl peptidase IV." JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 10, 1996, pages 2087-2093, XP002092856 usa the whole document	1,5

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

13 March 2000

Date of mailing of the international search report

24/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 6818 Patentkanal  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 681 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Kyriakakou, G



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 99/09708

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DOREEN M. ASHWORTH ET AL.: "2-Cyanopyrrolidides as potent, stable inhibitors of Dipeptidyl peptidase IV" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS,, vol. 6, no. 10, 1996, pages 1163-1166, XP004134889 the whole document	1,5
A	KJL AUGUSTYNS ET AL.: "Pyrrolidides: synthesis and structure-activity relationship as inhibitors of dipeptidyl peptidase IV" EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, vol. 32, 1997, pages 301-309, XP004086653 paris the whole document	1,5
A	CHEMICAL ABSTRACTS, vol. 123, no. 23, 4 December 1995 (1995-12-04) Columbus, Ohio, US; abstract no. 309272m, LI JINGRONG ET AL.: "Aminoacylpyrrolidine-2-nitriles: potent and stable inhibitors of dipeptidyl-peptidase IV." page 401; column 2; XP002132894 abstract & ARCH. BIOCHEM. BIOPHYS., vol. 323, no. 1, 1995, pages 148-154,	1,5

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/09708

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9819998 A	14-05-1998	AU 5318498 A	29-05-1998
		CN 1236361 A	24-11-1999
		CZ 9901615 A	11-08-1999
		EP 0937040 A	25-08-1999
		NO 992028 A	28-04-1999
		PL 332777 A	11-10-1999
WO 9515309 A	08-06-1995	AU 1113395 A	19-06-1995
		AU 8421998 A	12-11-1998
		CA 2178066 A	08-06-1995
		CN 1141033 A	22-01-1997
		CZ 9601595 A	15-01-1997
		EP 0731789 A	18-09-1996
		FI 962315 A	05-08-1996
		HU 76274 A	28-07-1997
		JP 9509921 T	07-10-1997
		NO 962269 A	30-07-1996
		PL 314838 A	30-09-1996
		US 5939560 A	17-08-1999
		ZA 9409525 A	02-08-1995